Medical Coverage Policy | Cochlear Implants



EFFECTIVE DATE: 05 | 17 | 2017

POLICY LAST UPDATED: 08 | 03 | 2022

OVERVIEW

A cochlear implant is a device for treatment of severe-to-profound hearing loss in individuals who only receive limited benefit from amplification with hearing aids. A cochlear implant provides direct electrical stimulation to the auditory nerve, bypassing the usual transducer cells that are absent or nonfunctional in deaf cochlea. This policy documents the coverage guidelines for cochlear implants.

MEDICAL CRITERIA

Not applicable

PRIOR AUTHORIZATION

Prior authorization is not required.

POLICY STATEMENT

Medicare Advantage Plans and Commercial Products

Cochlear Implantation

Unilateral or bilateral cochlear implantation of a U.S. Food and Drug Administration (FDA)-approved cochlear implant device may be considered medically necessary in patients age 9 months and older with bilateral severe to profound pre- or post-lingual (sensorineural) hearing loss and who have shown limited or no benefit from hearing aids.

Upgrades of an existing, functioning external system to achieve aesthetic improvement, such as smaller profile components or a switch from a body-worn, external sound processor to a behind-the-ear (BTE) model, are considered not covered, as this is considered a convenience. Additionally, replacement of internal and/or external components solely for the purpose of upgrading to a system with advanced technology or to a next-generation device is considered not covered, as this is considered a convenience.

Replacement of internal and/or external components is considered medically necessary only in a small subset of members who have inadequate response to existing component(s) to the point of interfering with the individual's activities of daily living, or the component(s) is/are no longer functional and cannot be repaired.

Hybrid Cochlear Implant/Hearing Aid

Cochlear implantation with a hybrid cochlear implant/hearing aid device that includes the hearing aid integrated into the external sound processor of the cochlear implant (eg, the Nucleus® HybridTM L24 Cochlear Implant System) may be considered medically necessary for patients ages 18 years and older who have bilateral severe-to-profound high-frequency sensorineural hearing loss with residual low frequency hearing sensitivity and receive limited benefit from appropriately fit bilateral hearing aids.

Medicare Advantage Plans

Cochlear implantation as a treatment for patients with unilateral hearing loss with or without tinnitus is not covered as the evidence is insufficient to determine the effects of the technology on health outcomes.

Commercial Products

Cochlear implantation as a treatment for patients with unilateral hearing loss with or without tinnitus is considered not medically necessary as the evidence is insufficient to determine the effects of the technology on health outcomes.

COVERAGE

Benefits may vary between groups/contracts. Please refer to the appropriate Benefit Booklet, Evidence of Coverage, or Subscriber Agreement for applicable Surgery Services and Medical Equipment, Medical Supplies, and Prosthetic Devices/Diagnostic Imaging, Lab, Machine Tests/Speech Therapy, and Personal Appearance and/or Items coverage/benefits.

BACKGROUND

A cochlear implant, classified by Centers for Medicare and Medicaid Services (CMS) as a prosthetic device, is an electronic instrument, part of which is implanted surgically to stimulate auditory nerve fibers, and part of which is worn or carried by the individual to capture, analyze, and code sound. Cochlear implant devices are available in single-channel and multi-channel models. The purpose of implanting the device is to provide awareness and identification of sounds and to facilitate communication for persons who are moderately to profoundly hearing impaired.

The basic structure of a cochlear implant includes both external and internal components. The external components include a microphone, an external sound processor, and an external transmitter. The internal components are surgically implanted and include an internal receiver implanted within the temporal bone and an electrode array that extends from the receiver into the cochlea through a surgically created opening in the round window of the middle ear.

Sounds that are picked up by the microphone are carried to the external sound processor, which transforms the sound into coded signals that are then transmitted through the skin to the implanted internal receiver. The receiver converts the incoming signals to electrical impulses that are then conveyed to the electrode array, ultimately resulting in stimulation of the auditory nerve.

A post-cochlear implant rehabilitation program is necessary to achieve benefit from the cochlear implant. A typical rehabilitation program consists of 6 to 10 sessions that last approximately $2^{1}/_{2}$ hours each. A rehabilitation program would include development of skills in understanding running speech, recognition of consonants and vowels, and tests of speech perception ability.

Typically, severe hearing loss is defined as a bilateral hearing threshold of 70 to 90 dB, and profound hearing loss is defined as a bilateral hearing threshold of 90 dB and above. In adults, limited benefit from hearing aids is defined as scores 50% correct or less in the ear to be implanted on tape-recorded sets of open-set sentence recognition. In children, limited benefit is defined as failure to develop basic auditory skills, and in older children, 30% or less correct on open-set tests.

Several cochlear implants are commercially available in the United States and are manufactured by Cochlear Americas, Advanced Bionics, and the MED-EL Corp. Over time, subsequent generations of the various components of the devices have been approved by the U.S. Food and Drug Administration (FDA), focusing on improved electrode design and speech-processing capabilities. Furthermore, smaller devices and the accumulating experience in children have resulted in broadening of the selection criteria to include children as young as 12 months.

In 2014, the Nucleus® Hybrid™ L24 Cochlear Implant System (Cochlear Americas) was approved by FDA through the premarket approval process. This system is a hybrid cochlear implant and hearing aid, with the hearing aid integrated into the external sound processor of the cochlear implant. It is indicated for unilateral use in patients ages 18 years and older who have residual low-frequency hearing sensitivity and severe-to-profound high-frequency sensorineural hearing loss, and who obtain limited benefit from an appropriately fit

bilateral hearing aid. The electrode array inserted into the cochlea is shorter than conventional cochlear implants. According to FDA's premarket approval notification, labeled indications for the device include:

- Preoperative hearing in the range from "normal to moderate hearing loss [HL] in the low frequencies (thresholds no poorer than 60 dB HL up to and including 500 Hz)"
- Preoperative hearing with "severe to profound mid to high frequency hearing loss (threshold average of 2000, 3000, and 4000 Hz ≥75 dB HL) in the ear to be implanted"
- Preoperative hearing with "moderately severe to profound mid to high frequency hearing loss (threshold average of 2000, 3000, and 4000 Hz ≥60 dB HL) in the contralateral ear"
- "The CNC [Consonant-Nucleus-Consonant] word recognition score will be between 10% and 60%, inclusively, in the ear to be implanted in the preoperative aided condition and in the contralateral ear equal to or better than that of the ear to be implanted but not more than 80% correct."

Other hybrid hearing devices have been developed. The Med-El EAS System received expanded PMA by the FDA in 2016 (PMA P000025/S084).

For individuals who have unilateral sensorineural hearing loss who receive the cochlear implant(s), the evidence for the use of cochlear implants in improving outcomes for patients with unilateral hearing loss, with or without tinnitus, is limited by small sample sizes, short follow-up times, and heterogeneity in evaluation protocols and outcome measurements. The evidence is insufficient to determine the effects of the technology on health outcomes.

CODING

In addition to the codes identified in this policy under the diagnostic imaging, lab, and machine tests benefit, there may be other therapeutic service codes related to cochlear implants (such as auditory rehabilitation) which would be applied to the member's speech therapy benefit.

Medicare Advantage Plans and Commercial Products

The following code is covered under the member's **surgery services** benefit:

69930 Cochlear device implantation, with or without mastoidectomy

The following codes are covered under the member's **speech therapy** benefit:

- 92626 Evaluation of auditory rehabilitation status; first hour
- 92627 Evaluation of auditory rehabilitation status; each additional 15 minutes
- 92630 Auditory rehabilitation; pre-lingual hearing loss
- 92633 Auditory rehabilitation; post-lingual hearing loss

The following codes are covered under the diagnostic imaging, lab, and machine tests benefit:

- 92601 Diagnostic analysis of cochlear implant, patient younger than 7 years of age; with programming
- **92602** Diagnostic analysis of cochlear implant, patient younger than 7 years of age; subsequent reprogramming
- 92603 Diagnostic analysis of cochlear implant, age 7 years or older; with programming
- 92604 Diagnostic analysis of cochlear implant, age 7 years or older; subsequent reprogramming

The following codes are covered under the **prosthetic devices** benefit:

- **L8614** Cochlear device, includes all internal and external components
- **L8615** Headset/headpiece for use with cochlear implant device, replacement
- **L8616** Microphone for use with cochlear implant device, replacement
- **L8617** Transmitting coil for use with cochlear implant device, replacement
- **L8618** Transmitter cable for use with cochlear implant device or auditory osseointegrated device, replacement
- L8619 Cochlear implant, external speech processor and controller, integrated system, replacement

- **L8621** Zinc air battery for use with cochlear implant device and auditory osseointegrated sound processors, replacement, each
- L8622 Alkaline battery for use with cochlear implant device, any size, replacement, each
- L8627 Cochlear implant, external speech processor, component, replacement
- L8628 Cochlear implant, external controller component, replacement
- L8629 Transmitting coil and cable, integrated, for use with cochlear implant device, replacement

The following codes are covered under the durable medical equipment benefit:

- **L8623** Lithium ion battery for use with cochlear implant device speech processor, other than ear level, replacement, each
- **L8624** Lithium ion battery for use with cochlear implant or auditory osseointegrated device speech processor, ear level, replacement, each
- **L8625** External recharging system for battery for use with cochlear implant or auditory osseointegrated device, replacement only, each

RELATED POLICIES

Clinical Trials Medicare Advantage Plans Clinical Trials Mandate Commercial Durable Medical Equipment Speech Therapy

PUBLISHED

Provider Update, October 2022 Provider Update, June 2021 Provider Update, September 2020 Provider Update, May 2020 Provider Update, September 2019

REFERENCES

- 1. Cochlear Implants in Adults and Children. NIH Consens Statement Online. 1995;13(2):1-30.
- 2. Bond M, Mealing S, Anderson R, et al. The effectiveness and cost-effectiveness of cochlear implants for severe to profound deafness in children and adults: a systematic review and economic model. Health Technol Assess. Sep 2009; 13(44): 1-330. PMID 19799825
- 3. Gaylor JM, Raman G, Chung M, et al. Cochlear implantation in adults: a systematic review and meta-analysis. JAMA Otolaryngol Head Neck Surg. Mar 2013; 139(3): 265-72. PMID 23429927
- 4. McRackan TR, Bauschard M, Hatch JL, et al. Meta-analysis of quality-of-life improvement after cochlear implantation and associations with speech recognition abilities. Laryngoscope. Apr 2018; 128(4): 982-990. PMID 28731538
- 5. McRackan TR, Bauschard M, Hatch JL, et al. Meta-analysis of Cochlear Implantation Outcomes Evaluated With General Health-related Patient-reported Outcome Measures. Otol Neurotol. Jan 2018; 39(1): 29-36. PMID 29227446
- 6. Crathorne L, Bond M, Cooper C, et al. A systematic review of the effectiveness and cost-effectiveness of bilateral multichannel cochlear implants in adults with severe-to-profound hearing loss. Clin Otolaryngol. Oct 2012; 37(5): 342-54. PMID 22928754
- 7. Choi JS, Betz J, Li L, et al. Association of Using Hearing Aids or Cochlear Implants With Changes in Depressive Symptoms in Older Adults. JAMA Otolaryngol Head Neck Surg. Jul 01 2016; 142(7): 652-7. PMID 27258813
- 8. van Zon A, Smulders YE, Ramakers GG, et al. Effect of unilateral and simultaneous bilateral cochlear implantation on tinnitus: A Prospective Study. Laryngoscope. Apr 2016; 126(4): 956-61. PMID 26255618
- 9. Bond M, Elston J, Mealing S, et al. Effectiveness of multi-channel unilateral cochlear implants for profoundly deaf children: a systematic review. Clin Otolaryngol. Jun 2009; 34(3): 199-211. PMID 19531168

- 10. Baron S, Blanchard M, Parodi M, et al. Sequential bilateral cochlear implants in children and adolescents: Outcomes and prognostic factors. Eur Ann Otorhinolaryngol Head Neck Dis. Apr 2019; 136(2): 69-73. PMID 30314876
- 11. Food and Drug Administration. Summary of Safety and Effectiveness Data (SSED): Nucleus 24 Cochlear Implant System (P970051/S172). 2020; https://www.accessdata.fda.gov/cdrh docs/pdf/P970051S172B.pdf. Accessed January 2, 2022.
- 12. Lyu J, Kong Y, Xu TQ, et al. Long-term follow-up of auditory performance and speech perception and effects of age on cochlear implantation in children with pre-lingual deafness. Chin Med J (Engl). Aug 20 2019; 132(16): 1925-1934. PMID 31365431
- 13. Karltorp E, Eklof M, Ostlund E, et al. Cochlear implants before 9 months of age led to more natural spoken language development without increased surgical risks. Acta Paediatr. Feb 2020; 109(2): 332-341. PMID 31350923
- 14. Sharma A, Dorman MF. Central auditory development in children with cochlear implants: clinical implications. Adv Otorhinolaryngol. 2006; 64: 66-88. PMID 16891837
- 15. Forli F, Arslan E, Bellelli S, et al. Systematic review of the literature on the clinical effectiveness of the cochlear implant procedure in paediatric patients. Acta Otorhinolaryngol Ital. Oct 2011; 31(5): 281-98. PMID 22287820
- 16. Sterkers F, Merklen F, Piron JP, et al. Outcomes after cochlear reimplantation in children. Int J Pediatr Otorhinolaryngol. Jun 2015; 79(6): 840-843. PMID 25843784
- 17. Black J, Hickson L, Black B, et al. Prognostic indicators in paediatric cochlear implant surgery: a systematic literature review. Cochlear Implants Int. May 2011; 12(2): 67-93. PMID 21756501
- 18. Pakdaman MN, Herrmann BS, Curtin HD, et al. Cochlear implantation in children with anomalous cochleovestibular anatomy: a systematic review. Otolaryngol Head Neck Surg. Feb 2012; 146(2): 180-90. PMID 22140206
- 19. Fernandes NF, Morettin M, Yamaguti EH, et al. Performance of hearing skills in children with auditory neuropathy spectrum disorder using cochlear implant: a systematic review. Braz J Otorhinolaryngol. Jan-Feb 2015; 81(1): 85-96. PMID 25458263
- 20. Vlastarakos PV, Proikas K, Papacharalampous G, et al. Cochlear implantation under the first year of agethe outcomes. A critical systematic review and meta-analysis. Int J Pediatr Otorhinolaryngol. Feb 2010; 74(2): 119-26. PMID 19896223
- 21. Ching TY, Dillon H, Day J, et al. Early language outcomes of children with cochlear implants: interim findings of the NAL study on longitudinal outcomes of children with hearing impairment. Cochlear Implants Int. 2009; 10 Suppl 1: 28-32. PMID 19067433
- 22. Colletti L, Mandala M, Zoccante L, et al. Infants versus older children fitted with cochlear implants: performance over 10 years. Int J Pediatr Otorhinolaryngol. Apr 2011; 75(4): 504-9. PMID 21277638
- 23. Guerzoni L, Murri A, Fabrizi E, et al. Social conversational skills development in early implanted children. Laryngoscope. Sep 2016; 126(9): 2098-105. PMID 26649815
- 24. Lammers MJ, van der Heijden GJ, Pourier VE, et al. Bilateral cochlear implantation in children: a systematic review and best-evidence synthesis. Laryngoscope. Jul 2014; 124(7): 1694-9. PMID 24390811
- 25. Broomfield SJ, Murphy J, Emmett S, et al. Results of a prospective surgical audit of bilateral paediatric cochlear implantation in the UK. Cochlear Implants Int. Nov 2013; 14 Suppl 4: S19-21. PMID 24533758
- 26. Sarant J, Harris D, Bennet L, et al. Bilateral versus unilateral cochlear implants in children: a study of spoken language outcomes. Ear Hear. Jul-Aug 2014; 35(4): 396-409. PMID 24557003
- 27. Escorihuela Garcia V, Pitarch Ribas MI, Llopez Carratala I, et al. Comparative study between unilateral and bilateral cochlear implantation in children of 1 and 2 years of age. Acta Otorrinolaringol Esp. May-Jun 2016; 67(3): 148-55. PMID 26632253
- 28. Friedmann DR, Green J, Fang Y, et al. Sequential bilateral cochlear implantation in the adolescent population. Laryngoscope. Aug 2015; 125(8): 1952-8. PMID 25946482
- 29. Illg A, Giourgas A, Kral A, et al. Speech comprehension in children and adolescents after sequential bilateral cochlear implantation with long interimplant interval. Otol Neurotol. Jun 2013; 34(4): 682-9. PMID 23640090

- 30. van Zon A, Peters JP, Stegeman I, et al. Cochlear implantation for patients with single-sided deafness or asymmetrical hearing loss: a systematic review of the evidence. Otol Neurotol. Feb 2015; 36(2): 209-19. PMID 25502451
- 31. Benchetrit L, Ronner EA, Anne S, et al. Cochlear Implantation in Children With Single-Sided Deafness: A Systematic Review and Meta-analysis. JAMA Otolaryngol Head Neck Surg. Jan 01 2021; 147(1): 58-69. PMID 33151295
- 32. Marx M, Mosnier I, Venail F, et al. Cochlear Implantation and Other Treatments in Single-Sided Deafness and Asymmetric Hearing Loss: Results of a National Multicenter Study Including a Randomized Controlled Trial. AudiolNeurootol. 2021; 26(6): 414-424. PMID 33789270
- 33. Peters JPM, van Heteren JAA, Wendrich AW, et al. Short-term outcomes of cochlear implantation for single-sideddeafness compared to bone conduction devices and contralateral routing of sound hearing aids-Results of aRandomised controlled trial (CINGLE-trial). PLoS One. 2021; 16(10): e0257447. PMID 34644322
- 34. Buss E, Dillon MT, Rooth MA, et al. Effects of Cochlear Implantation on Binaural Hearing in Adults With Unilateral Hearing Loss. Trends Hear. Jan-Dec 2018; 22: 2331216518771173. PMID 29732951
- 35. Dillon MT, Buss E, O'Connell BP, et al. Low-Frequency Hearing Preservation With Long Electrode Arrays: Inclusion of Unaided Hearing Threshold Assessment in the Postoperative Test Battery. Am J Audiol. Mar 05 2020; 29(1): 1-5. PMID 31835906
- 36. Galvin JJ, Fu QJ, Wilkinson EP, et al. Benefits of Cochlear Implantation for Single-Sided Deafness: Data From the House Clinic-University of Southern California-University of California, Los Angeles Clinical Trial. Ear Hear. Jul/Aug 2019; 40(4): 766-781. PMID 30358655
- 37. Peter N, Kleinjung T, Probst R, et al. Cochlear implants in single-sided deafness clinical results of a Swiss multicentre study. Swiss Med Wkly. Dec 16 2019; 149: w20171. PMID 31880806
- 38. Poncet-Wallet C, Mamelle E, Godey B, et al. Prospective Multicentric Follow-up Study of Cochlear Implantation in Adults With Single-Sided Deafness: Tinnitus and Audiological Outcomes. Otol Neurotol. Dec 20 2019. PMID 31868784
- 39. Dillon MT, Buss E, Rooth MA, et al. Cochlear Implantation in Cases of Asymmetric Hearing Loss: Subjective Benefit, Word Recognition, and Spatial Hearing. Trends Hear. Jan-Dec 2020; 24: 2331216520945524. PMID 32808881
- 40. Food and Drug Administration. Post-Approval Studies (PAS): MED-EL New Enrollment SSD/AHL Study. 2020;
 - https://www.accessdata.fda.gov/scripts/cdrh/cfdocs/cfpma/pma_pas.cfm?t_id=647845&c_id=5585. Accessed on January 4, 2022.
- 41. Food and Drug Administration. Summary of Safety and Effectiveness Data (SSED): MED-EL Cochlear Implant System (P000025/S104). 2019; https://www.accessdata.fda.gov/cdrh_docs/pdf/P000025S104B.pdf. Accessed January 16, 2021.
- 42. Mertens G, De Bodt M, Van de Heyning P. Cochlear implantation as a long-term treatment for ipsilateral incapacitating tinnitus in subjects with unilateral hearing loss up to 10 years. Hear Res. Jan 2016; 331: 1-6. PMID 26433053
- 43. Rahne T, Plontke SK. Functional Result After Cochlear Implantation in Children and Adults With Single-sided Deafness. Otol Neurotol. Oct 2016; 37(9): e332-40. PMID 27631656
- 44. Vlastarakos PV, Nazos K, Tavoulari EF, et al. Cochlear implantation for single-sided deafness: the outcomes. An evidence-based approach. Eur Arch Otorhinolaryngol. Aug 2014; 271(8): 2119-26. PMID 24096818
- 45. Ramos Macias A, Falcon Gonzalez JC, Manrique M, et al. Cochlear implants as a treatment option for unilateral hearing loss, severe tinnitus and hyperacusis. Audiol Neurootol. 2015; 20 Suppl 1: 60-6. PMID 25997672
- 46. Tavora-Vieira D, Marino R, Krishnaswamy J, et al. Cochlear implantation for unilateral deafness with and without tinnitus: a case series. Laryngoscope. May 2013; 123(5): 1251-5. PMID 23553411
- 47. Pillsbury HC, Dillon MT, Buchman CA, et al. Multicenter US Clinical Trial With an Electric-Acoustic Stimulation (EAS) System in Adults: Final Outcomes. Otol Neurotol. Mar 2018; 39(3): 299-305. PMID 29342054

- 48. Food and Drug Administration. Approval Letter: Nucleus Hybrid L24 Cochlear Implant System (P130016). 2014; https://www.accessdata.fda.gov/cdrh_docs/pdf13/P130016a.pdf. Accessed January 5, 2022.
- 49. Roland JT, Gantz BJ, Waltzman SB, et al. United States multicenter clinical trial of the cochlear nucleus hybrid implant system. Laryngoscope. Jan 2016; 126(1): 175-81. PMID 26152811
- 50. Roland JT, Gantz BJ, Waltzman SB, et al. Long-term outcomes of cochlear implantation in patients with high-frequency hearing loss. Laryngoscope. Aug 2018; 128(8): 1939-1945. PMID 29330858
- 51. Lenarz T, James C, Cuda D, et al. European multi-centre study of the Nucleus Hybrid L24 cochlear implant. Int J Audiol. Dec 2013; 52(12): 838-48. PMID 23992489
- 52. Santa Maria PL, Gluth MB, Yuan Y, et al. Hearing preservation surgery for cochlear implantation: a meta-analysis. Otol Neurotol. Dec 2014; 35(10): e256-69. PMID 25233333
- 53. Causon A, Verschuur C, Newman TA. A Retrospective Analysis of the Contribution of Reported Factors in Cochlear Implantation on Hearing Preservation Outcomes. Otol Neurotol. Aug 2015; 36(7): 1137-45. PMID 25853614
- 54. Gantz BJ, Dunn C, Oleson J, et al. Multicenter clinical trial of the Nucleus Hybrid S8 cochlear implant: Final outcomes. Laryngoscope. Apr 2016; 126(4): 962-73. PMID 26756395
- 55. American Academy of Otolaryngology -- Head and Neck Surgery. Position Statement: Cochlear Implants. November 10, 2020; http://www.entnet.org/Practice/policyCochlearImplants.cfm. Accessed January 7, 2022.
- 56. Raman G, Lee J, Chung MG, et al. Technology Assessment Report: Effectiveness of Cochlear Implants in Adults with Sensorineural Hearing Loss Rockville, MD: Agency for Healthcare Research and Quality; 2011
- 57. National Institute for Health and Care Excellence (NICE). Cochlear Implants for Children and Adults With Severe to Profound Deafness [TA566]. 2019; https://www.nice.org.uk/guidance/ta566/. Accessed January 7, 2022.
- 58. Centers for Medicare & Medicaid. Cochlear Implantation. 2013; https://www.cms.gov/Medicare/Coverage/Coverage-with-Evidence-Development/Cochlear-Implantation-.html. Accessed January 7, 2022.

----- CLICK THE ENVELOPE ICON BELOW TO SUBMIT COMMENTS

This medical policy is made available to you for informational purposes only. It is not a guarantee of payment or a substitute for your medical judgment in the treatment of your patients. Benefits and eligibility are determined by the member's subscriber agreement or member certificate and/or the employer agreement, and those documents will supersede the provisions of this medical policy. For information on member-specific benefits, call the provider call center. If you provide services to a member which are determined to not be medically necessary (or in some cases medically necessary services which are non-covered benefits), you may not charge the member for the services unless you have informed the member and they have agreed in writing in advance to continue with the treatment at their own expense. Please refer to your participation agreement(s) for the applicable provisions. This policy is current at the time of publication; however, medical practices, technology, and knowledge are constantly changing. BCBSRI reserves the right to review and revise this policy for any reason and at any time, with or without notice. Blue Cross & Blue Shield of Rhode Island is an independent licensee of the Blue Cross and Blue Shield Association.

